

MEMORANDUM

To: School Nurses

From: Linda Ionta, Director of Athletics,
Health, Physical Education & Nurses

Re: School Health Standing Orders

Date: September 2016

Attached, please find the updated School Health Standing Orders to replace into the Medical Service Handbook.

If you have any concerns/questions, please contact me.

Copy to: (w/att.)
Diane Cappiello – Business Office

UNION TOWNSHIP PUBLIC SCHOOLS
Union, New Jersey 07083

PHYSICIAN'S STANDING ORDERS FOR PRN MEDICATION


The following medication may be administered by the School Nurse to Board of Education employees and to students at Union High School, Burnet and Kawameeh Middle Schools, who weigh more than 96 pounds, for the purpose of treating mild symptoms. Students must also have parental or legal guardian's consent for these medications to be administered. If symptoms persist it is understood that the nurse will contact the parent or guardian.

Acetaminophen

325 mg, 2 tablets orally, every 4 hours as needed for pain or fever.

Ibuprofen

200 mg, 1 or 2 tablets orally, every 6 hours as needed for pain or menstrual cramps.



Bharati S. Mullick, M.D., F.A.A.P.
Chief School Medical Examiner

8/23/16

Date

**TOWNSHIP OF UNION PUBLIC SCHOOLS
Union, NJ 07083**

PHYSICIAN'S STANDING ORDERS FOR ROUTINE MEDICATION

The following pharmaceuticals, treatments, and tests may be administered per package directions as needed by the Certified School Nurse or Registered Nurse Substitute to students and school board employees:

Anbesol topically
Bactine topically
Bacitracin topically
Calamine lotion topically
Hibiclens Skin and Wound Cleanser topically
Hydrocortisone 1 % topically
Hydrogen Peroxide topically
Insect Sting Swab topically
Isopropyl Alcohol topically
Petroleum Jelly topically
Witch Hazel topically


Cepacol Throat Lozenges orally (adults and children ages 6 and older)
Chloraseptic Throat Gargle or Spray orally for ages 6 years and older
Gyloxide Oral Antiseptic, swish and spit

Dacriose Ophthalmic Irrigation
Visine Eye Drops for allergy relief


T.B. Testing: Mantoux Tuberculin, PPD (5 TU/0.1mL)
Administer 0.1mL intradermally, read in 48 to 72 hours

Emergency Standing Orders:

Benadryl (see attached)
Epinephrine (see attached)



Bharati S. Mullick, M.D., F.A.A.P.
Chief School Medical Examiner



Date

UNION TOWNSHIP PUBLIC SCHOOLS
Union, New Jersey 07083

PHYSICIAN'S STANDING ORDERS FOR EMERGENCY MEDICATION

Benadryl (Diphenhydramine HCl) may be administered to students and Board of Education employees by the School Nurse in the following emergency situation:

For acute allergic reactions including hay fever, insect stings, etc.

Dosage: Administer Benadryl Elixir 12.5 mg/tsp. or
Benadryl 25 mg capsules based on student's weight

If student weighs 30 to 45 pounds, administer 1 to 2
teaspoons orally


If student weighs 46 to 90 pounds, administer 2 to 4
teaspoons or 1 to 2 capsules orally

If student weighs more than 90 pounds, administer 1 to 2
capsules orally


Contraindications: Known hypersensitivity to Benadryl

Side Effects: Drowsiness, dry mouth, pallor, anxiety. See package directions
and alerts.

Nursing Implications: Monitor student's allergic symptoms and response to medication.



Bharati S. Mullick, M.D., F.A.A.P.
Chief School Medical Examiner



Date

UNION TOWNSHIP PUBLIC SCHOOLS

Union, New Jersey 07083

COMMUNICABLE DISEASE POLICIES

1. ***Chicken Pox***

Usually begins with fever, fatigue, and a skin eruption that starts as a red papule, then becomes small blisters for 3-4 days, then leaves a scab. This disease is communicable from 24 hours before to 6-7 days after the eruption at which time the blisters have dried up and become scabbed.

- . The student will be admitted to class seven (7) days after the rash appears if all lesions are dry.

2. ***Conjunctivitis (Pink Eye)***

Defined as pink or red conjunctiva with white or yellow discharge, often with matted eyelids after sleep and eye pain or redness of the eyelids or skin surrounding the eye.

- . The student may return to school after examination by a physician and approved for readmission, with or without treatment.

3. ***Strep Throat Infections***

Usually characterized by a sore throat for four days with no fever, or a sore throat and fever for two days.

- . The student may return to school 24 hours after antibiotic treatment has begun and 24 hours after fever has subsided.

4. ***Pediculosis Capitis (Head Lice)***

The student will be excluded from school until no nits are visible.

- . Recheck head after returning to school.

5. ***Tinea Corporis (Ringworm)***

Usually begins as tiny red spot, which slowly grows in circular fashion, clearing in the center as it enlarges. The edges remain reddish and scaly.


- . Student must have medical note to return to school
- . Exposed skin must be kept covered

6. ***Tinea Pedis (Athlete's Foot)***

Characterized by peeling, cracking and itching of skin between the toes. This is a superficial infection widely spread through the use of showers, swimming pools, common bath mats or direct contact with an infected person. Isolation is not practical.

7. ***Skin Eruptions (Impetigo, Staph)***

The student should be excluded from school until satisfactory treatment has been started and a note is received from a physician that the student has permission to return. The student should be excused from Physical Education and showers until lesions are completely healed.


Bharati S. Mullick, MD, F.A.A.P.
Chief School Medical Examiner

8/23/16
Date

UNION TOWNSHIP PUBLIC SCHOOLS
Union, New Jersey 07083

PHYSICIAN'S STANDING ORDERS FOR EMERGENCY MEDICATION

Epinephrine (adrenalin) may be administered by the School Nurse in the following emergency situations:

1. Anaphylactic shock
2. Respiratory failure or severe laryngeal edema due to severe reaction to insect bites or stings.

Dosage: Adult Epinephrine Auto-Injector 0.3mg for students weighing more than 66 pounds

Pediatric Epinephrine Auto-Injector 0.15mg for students weighing 66 pounds or less

Route: Intramuscular in anteriolateral aspect of the thigh, may be repeated in 15 minutes.


Contraindications: Known previous hypersensitivity to Epinephrine.

Side Effects: Nervousness, tremor, palpitation, anxiety, headache, or elevated blood pressure.

Nursing Implications: Do not use a brown or precipitated solution.
Monitor vital signs as indicated.
Injection site should be massaged after administration of Epinephrine.
Transport student to emergency room ASAP.



Bharati S. Mullick, M.D., F.A.A.P.
Chief School Medical Examiner



Date